

Pain Control Surrounding Castration and Tail Docking in Lambs

Tail docking and castration are important management procedures for the sheep industry globally; however, they are both painful procedures that can impact animal welfare. Therefore, it is critical that these procedures are performed using best management practices, including pain control.

Take Home Messages

- Tail docking and castration are painful no matter the method that is applied, with rises in cortisol and pain behaviors noted following the procedures
- From available literature, it seems there are differences in the level of pain experienced when different methods are used, with the use of a rubber ring causing the highest level of pain for both castration and tail docking.
- To control pain, the combination of a local anesthetic and NSAID may result in the best pain relief for both tail docking and castration

Tail Docking

Tail docking is commonly practiced to manage fecal soiling to reduce the risk of fly strike (myiasis) (French et al., 1994) and is completed using many different methods. No matter the method, it is painful as it stimulates nociceptors by causing tissue damage and by triggering physiological pain pathways (Meintjes, 2012). As a result of the pain associated with tail docking, lambs will commonly exhibit behaviors, such as vocalization, tail wagging, and restlessness.

Does Method Matter for Pain?

From a study by Grant (2004), it was found that it does when the use of tail docking by hot iron or by rubber ring was directly compared.

Hot Iron Docking

When evaluating hot docking, the largest pain response was seen at the time of application with vocalizing and struggling commonly observed. In addition, docking irons that had the temperature too low resulted in slow and prolonged cutting times, whereas high docking iron temperatures led to rapid cutting but inadequate cauterization, highlighting the need to have properly maintained and functioning equipment. Other than the initial pain response, few indicators of pain occurred other than an increase in active pain behaviors observed 40 min following the tail docking. This response is consistent with burn pain where shock and destruction of nerve endings from deep injuries delay pain.

Rubber Ring Docking

Tail docking with rubber rings lead to large behavioral and postural responses associated with pain. The response was much greater than hot iron docking. The reason for this is that the tight rubber rings prevent blood supply to the tail but don't prevent the conduction of nerve impulses from the painful, ischemic tail (Molony et al., 1993; Kent et al., 1995). In contrast, with hot docking, the nociceptors at the tail stump are destroyed or the removal of the entire tail leads to all input from nociceptive receptors contained within the tail to be eliminated, leaving only receptors in the stump.

What About Other Methods?

Both Molony et al. (1993) and Kent et al. (1995) compared other methods to rubber ring docking. Specifically, Molony et al. (1993) compared the use of a Burdizzo, surgery or rubber ring docking and found that the use of a Burdizzo may be the least painful method, with the rubber ring being most painful. Similarly, Kent et al. (1995) found that the Burdizzo was a less painful method.

Mitigating Pain Associated With the Different Methods of Tail Docking

Rubber Ring Docking

Using a local anesthetic administered subcutaneously at the site of application or giving an epidural with a local anesthetic prior to the application of the ring has been shown to reduce cortisol responses and pain behavior in the hours following application (Kells et al., 2020; Graham et al., 1997; Kent et al., 1998).

Beyond the application of local anesthetic, a new product is currently under development that has lidocaine implanted directly in the band (Saville et al, 2020). This may allow for the slow release of lidocaine into the site of application over time.

With regard to non-steroidal anti-inflammatory drugs, several studies have shown a benefit. Graham et al. (1997) found that the injection of an NSAID reduced cortisol response but no behavioral indicators of pain, whereas Pollard et al. (2001) found the oral administration of an NSAID reduced restlessness, active behavior, and abnormal posture following tail docking.

Other Methods

Similar to rubber ring castration, benefits have been shown to providing a local anesthetic, specifically when hot docking and completing surgical removal (Dunthorne, 2021). For NSAID administration, further evidence supports its use, especially when hot docking. In lambs that were either provided with meloxicam or a placebo at a combined hot docking of the tail and surgical castration, a large reduction was found in behaviors indicative of pain as well as an increase in the time spent grazing, suckling, and lying was observed in the meloxicam group (Small et al., 2014). Kells et al. (2019) found that using meloxicam in combination with a local anesthetic with hot docking and rubber ring castration led to reduced behavioral signs of pain.

Cumulatively, there is an existing body of evidence highlighting that to best manage pain a combination of a local anesthetic with an NSAID is required (Dunthorne, 2021).

Castration

Castration is commonly practiced in ram lambs to eliminate sexual behavior and reduce fighting as well as allow male and female lambs to be managed together. In addition, there is no risk of ram taint in rams that have been castrated which improves meat quality. There are few alternatives to performing castration and it is a necessary practice in the industry.

Does Method Matter for Pain?

The method of castration also influences the level of pain that is noted following castration.

Surgical Castration

Surgical castration causes the largest spike in cortisol in the hours following castration compared to other methods (Mellor and Stafford, 2000). When compared to a clamp method, an increased cortisol response was noted in the first 6 or 9 hours following the procedure as well as an increased amount of time with an abnormal posture and a larger response to scrotal palpation (Melches et al., 2007). This highlights the need to control pain, especially acute pain, when applying this method.

Rubber Ring Castration

Similar to tail docking, rubber ring castration could be one of the more painful methods of castration. Compared to the use of a clamp (i.e., Burdizzo), ring castration increased abnormal postures (Dinniss et al., 1999), whereas compared to surgical castration, there were higher levels of abnormal lying behavior and restlessness; although, lower cortisol was noted (Colditz et al., 2012; Lester et al., 1991). In addition, until the scrotum falls off after 4 weeks following the application of the rubber ring, it remains swollen and appears to cause behavioral signs of discomfort (Cockram et al., 2012).

Clamp Castration

Using a clamp technique (i.e., Burdizzo) has been shown to cause an increased level of cortisol and for the first 15 minutes after the procedure an increased number of pain behaviours are shown (Mellor et al., 1991). However, when compared to other techniques, the signs of pain are found to be lower as highlighted above. Interestingly, using a clamp prior to the placement of a rubber ring seems to decrease some pain associated with ring castration alone (Kent et al., 1998; Thorton and Waterman-Pearson, 1999); however, it is more painful than compared to solely using a clamp (Molony et al., 1997).

Controlling Pain Associated with Castration

Surgical Castration

Local anesthetic injected into the scrotal neck, cord, and testis can reduce plasma cortisol in response to surgical castration (Thorton and Waterman-Pearson, 1999); however, an increased cortisol level is seen 3 to 6 hours following the administration of a local anesthetic (Melches et al., 2007). Although, there have not been any formal studies evaluating the use of a local anesthetic in combination with an NSAID for surgical castration, it would seem that solely a local anesthetic on its own may not be enough to achieve optimal pain control.

Rubber Ring Castration

Administration of local anesthetic into the scrotal neck and cord or the testis can lead to reduced cortisol responses as well as behavioral indicators of pain and discomfort, especially in the period immediately following application of the band (Kent et al., 1998; Thornton and Waterman-Pearson, 1999).

When it comes to the use of NSAIDs, few studies have been completed. However, one study found that the injection of an NSAID reduced cortisol and behavioral responses associated with pain in the hours following castration (Paull et al., 2012). Others have found few differences on acute pain behaviour (Kells et al., 2020) or a slightly reduced ADG but lower level of lamb losses (Small et al., 2020). More research is needed to highlight the utility of using NSAIDs to control chronic pain.

Clamp Castration

Using a combination of a local anesthetic and NSAID likely provides optimal pain control when the clamp technique is used. Durand et al. (2019) found that using a local anesthetic (administered into each spermatic cord and around the scrotal neck) and an NSAID, flunixin, lead to reduced behavioral signs of pain and discomfort and lower cortisol levels, with the authors suggesting multimodal analgesia for up to 3 days post castration.

References

- Cockram, M.S., P. Menzies, D. Barrett, and J. Hemsted. 2012. Code of practice for the care and handling of sheep: Review of scientific research on priority issues.
- Colditz, I., D. Paull, and C. Lee. 2012. Social transmission of physiological and behavioral responses to castration in suckling Merino lambs. *Appl Anim Behav Sci.* 136:136-145.
- Dinniss, A., K. Stafford, D. Mellor, R. Bruce, and R. Ward. 1999. The behaviour pattern of lambs after castration using a rubber ring and/or castrating clamp with or without local anesthetic. *New Zealand Vet J.* 47:198-203.
- Dunthorne, E. 2021. Tail docking and castrating lambs: does the administration of local anesthetic or meloxicam reduce the pain response exhibited? *Veterinary Evidence.*
- Durand, D., M. Faure, A. de la Foye, and A. De Boyer des Roches. 2019. Benefits of a multimodal analgesics compared to local anesthesia alone to alleviate pain following castration in sheep: a multiparametric approach. *Animal.*
- Grant, C. 2004. Behavioural responses of lambs to common painful husbandry procedures. *Applied Animal Behaviour Science.* 87:255-273.
- French, N.P., R. Wall, and K.L. Morgan. 1994. Lamb tail docking: a controlled field study of the effects of tail amputation on health and productivity. *Vet Rec.* 134: 463.
- Kells, N.J., Beausoleil, N.J., Godfrey, A.J.R., Littlewood, K.E., Ward, R.N. & Johnson, C.B. (2019). Effect of analgesic strategies on pain behaviour associated with combined ring castration and hot iron tail docking in Merino lambs. *Applied Animal Behaviour Science.* 222.
- Kent, J., V. Molony, and I. Robertson. 1995. Comparison of the burdizzo and rubber ring methods for castrating and tail docking lambs. *Vet Rec.* 136:192-196.
- Kent, J., V. Molony, and M. Graham. 1998. Comparison of methods for the reduction of acute pain produced by rubber ring castration or tail docking week-old lambs. *Vet J.* 155: 39-51.
- Lester, S., D. Mellor, R. Ward, and R. Holmes. 1991. Cortisol responses to young lambs to castration and tailing using different methods. *New Zealand Vet J.* 39:134-138.
- Meintjes, R.A. 2012. An overview of the physiology of pain for the veterinarian. *Vet J.* 193:344-348.
- Melches, S., S.C. Mellema, M.G. Doherr, B. Wechsler, and A. Steiner. 2007. Castration of lambs: A welfare comparison of different castration techniques in lambs over 10 weeks of age. *Vet J.* 173:554-563.
- Mellor, D.J., V. Molony, and I.S. Robertson. 1991. Effects of castration on behaviour and plasma-cortisol concentrations in young lambs, kids, and calves. *Res Vet Sci.* 51:149-154.
- Mellor, D.J., and K.J. Stafford. 2000. Acute castration and/or tailing distress and its alleviation in lambs. *New Zealand Vet J.* 48:33-43.
- Molony, V., J. Kent, and I. Robertson. 1993. Behavioral responses of lambs of three ages in the first three hours after three methods of castration and tail docking. *Res. Vet Sci.* 55: 236-245.
- Molony, V., J.E. Kent, B.D. Hosie, and M.J. Graham. 1997. Reduction in pain suffered by lambs at castration. *Vet J.* 153:205-213.
- Pollard, J.C., V. Roos, and R.P. Littlejohn. 2001. Effects of an oral dose of acetyl salicylate at tail docking on the behaviour of lambs aged three to six weeks. *Appl Anim Behav Sci.* 71:29-42.

Saville, J. W., Ross, J. A., Trefz, T., Schatz, C., Matheson-Bird, H., Ralston, B., Granot, O., Schmid, K., Terry, R., **Allan, N.D.**, Wulff, J.E., Olson, M. (2020). Development and Field Validation of Lidocaine-Loaded Castration Bands for Bovine Pain Mitigation. *Animals*, 10(12), 1-16. <https://doi.org/10.3390/ani10122363>.

Small, A., Belson, S., Holm, M. & Colditz, I.G. (2014). Efficacy of a buccal meloxicam formulation for pain relief in Merino lambs undergoing knife castration and tail docking in a randomized field trial. *Australian Veterinary Journal*. 92(10), 318-388.

Small, A.H., S. Belson, H. Brewer, and S.M. Schmoelzl. 2020. Marking to weaning production aspects of lambs provided with NSAID analgesia compared with lambs receiving no analgesia at the time of elastrator ring marking. *Aust Vet J*. 99:40-43.

Thorton, P.D., and A.E. Waterman-Pearson. 1999. Quantification of the pain and distress responses to castration in young lambs. *Res Vet Sci*. 66:107-118.

